

Cosmeceutical Role of Medicinal Plants and Herbs: Therapeutic and Cosmetic Perspectives

Zarin Fatima¹, Simran Khan², Heena Khan¹, Shahla Khanam³ and Ahmad Zia^{*4}

¹Department of Saidla, Ajmal Khan Tibbiya College, Aligarh Muslim University, Aligarh, UP, India

²Department of Mahiyatul Amraz (Pathology), Government Tibbi College and Hospital, Patna, Bihar, India

³Department of Tashreeh wa Manafeul Aza, Ajmal Khan Tibbiya College, Aligarh Muslim University, Aligarh, UP, India

⁴Department of Tashreeh-ul-Badan, State Takmeel-ut-Tib College and Hospital, Lucknow, Uttar Pradesh, India

Citation: Zarin Fatima, Simran Khan, Heena Khan, Shahla Khanam, and Ahmad Zia (2026). Cosmeceutical Role of Medicinal Plants and Herbs: Therapeutic and Cosmetic Perspectives. *Acta Traditional Medicine*.

DOI: <https://doi.org/10.51470/ATM.2026.5.1.16>

Corresponding Author: **Ahmad Zia** | E-Mail: ziaahmad3000@gmail.com

Received 10 October 2025 | Revised 11 November 2025 | Accepted 11 December 2025 | Available Online January 09 2026

ABSTRACT

Cosmeceuticals represent a convergence of cosmetics and pharmaceuticals, incorporating bioactive compounds from medicinal plants that offer both therapeutic and aesthetic benefits. This review explores the multifaceted roles of plant-derived ingredients in skincare, examining their anti-inflammatory, antioxidant, antimicrobial, anti-aging, and skin-regenerative properties. Important phytochemicals including polyphenols, terpenoids, alkaloids, and vitamins demonstrate efficacy in addressing photoaging, hyperpigmentation, acne, and dermatological disorders through diverse molecular mechanisms. Notable botanical sources such as green tea, turmeric, aloe vera, and ginseng provide evidence-based alternatives to synthetic ingredients. Despite promising therapeutic potential, challenges persist regarding standardization, bioavailability, clinical validation, and regulatory frameworks. Advanced extraction technologies and novel delivery systems continue to enhance phytocosmeceutical efficacy. This comprehensive analysis bridges traditional botanical knowledge with contemporary scientific evidence, elucidating the future trajectory of plant-based cosmeceutical development and evidence-based skincare innovation.

Keywords: Cosmeceuticals; Medicinal plants; Phytochemicals; Skin aging; Herbal cosmetics; Natural antioxidants.

Introduction

The intersection of cosmetics and pharmaceuticals has given rise to the rapidly evolving field of cosmeceuticals, where bioactive ingredients derived from medicinal plants and herbs serve both therapeutic and aesthetic purposes. Cosmeceuticals represent a unique category of topical products that transcend traditional cosmetics by incorporating biologically active compounds capable of influencing skin structure and function at the cellular level [1]. Unlike conventional cosmetics that merely enhance appearance temporarily, cosmeceuticals claim to deliver tangible physiological benefits through their pharmacologically active constituents. Medicinal plants have been integral to skincare and beauty regimens across diverse cultures for millennia, with ancient civilizations such as Egyptian, Ayurvedic, and Traditional Chinese Medicine systems documenting extensive use of botanical preparations for dermatological applications [2]. The modern resurgence of interest in plant-based cosmeceuticals stems from mounting concerns about synthetic chemical ingredients, growing consumer demand for natural and sustainable products, and accumulating scientific evidence supporting the efficacy of phytochemical compounds [3]. Contemporary research has validated many traditional uses of medicinal herbs, revealing that plants produce a remarkable array of secondary metabolites—including polyphenols, terpenoids, alkaloids, and flavonoids—that exhibit potent antioxidant, anti-inflammatory, antimicrobial, and anti-aging properties.

The skin, as the largest organ of the human body, is continuously exposed to environmental stressors such as ultraviolet radiation, pollution, and oxidative stress, which accelerate aging processes and contribute to various dermatological disorders [4]. Plant-derived cosmeceuticals offer multifaceted protection through diverse mechanisms: neutralizing free radicals, inhibiting matrix metalloproteinases, stimulating collagen synthesis, reducing hyperpigmentation, and maintaining skin barrier integrity [5]. Notable examples include green tea (*Camellia sinensis*) polyphenols for photoprotection, turmeric (*Curcuma longa*) curcuminoids for anti-inflammatory effects, aloe vera (*Aloe barbadensis*) polysaccharides for wound healing, and ginseng (*Panax ginseng*) ginsenosides for anti-wrinkle activity [6]. Despite the promising therapeutic potential of plant-based cosmeceuticals, several challenges persist in their development and commercialization. Issues related to standardization of botanical extracts, variability in phytochemical composition, stability concerns, bioavailability limitations, and the need for rigorous clinical validation remain critical areas requiring attention [7]. Furthermore, regulatory frameworks governing cosmeceuticals vary significantly across different countries, creating complexities in product classification and marketing claims. This comprehensive review explores the cosmeceutical role of medicinal plants and herbs from both therapeutic and cosmetic perspectives, examining the scientific basis for their dermatological applications, mechanisms of action of key phytoactive compounds, current market trends, regulatory considerations, and future directions

in phytocosmeceutical research. The bridging traditional botanical wisdom with modern scientific investigation, this discussion aims to elucidate how plant-derived ingredients continue to shape the future of evidence-based skincare and aesthetic medicine.

Therapeutic Roles of Medicinal Plants in Cosmeceuticals

Anti-inflammatory and Healing Properties

Medicinal plants have demonstrated remarkable anti-inflammatory and wound-healing capabilities through various phytochemical constituents. Aloe vera (*Aloe barbadensis*) contains polysaccharides, particularly acemannan, which accelerate wound healing by stimulating fibroblast proliferation and collagen synthesis [8]. Calendula (*Calendula officinalis*) exhibits potent anti-inflammatory effects through triterpene esters that reduce cytokine production and promote tissue regeneration [9]. Chamomile (*Matricaria chamomilla*) contains α -bisabolol and chamazulene, which inhibit prostaglandin synthesis and mitigate inflammatory skin conditions including eczema and dermatitis.

Antioxidant Effects in Skin Protection

Plant-derived antioxidants neutralize reactive oxygen species (ROS) that cause oxidative damage to cellular components, thereby preventing premature aging and photoaging. Green tea (*Camellia sinensis*) polyphenols, particularly epigallocatechin-3-gallate (EGCG), demonstrate superior free radical scavenging activity and protect against UV-induced DNA damage [10]. Grape seed (*Vitis vinifera*) proanthocyanidins exhibit strong antioxidant properties, protecting dermal fibroblasts from oxidative stress and maintaining skin elasticity [11]. Pomegranate (*Punica granatum*) ellagitannins provide photoprotection by inhibiting matrix metalloproteinases and preserving collagen integrity.

Antimicrobial and Antifungal Activities

Numerous medicinal plants possess broad-spectrum antimicrobial properties valuable in treating skin infections and maintaining skin microbiome balance. Tea tree oil (*Melaleuca alternifolia*) contains terpinen-4-ol, demonstrating efficacy against *Propionibacterium acnes*, *Staphylococcus aureus*, and various fungal pathogens [12]. Neem (*Azadirachta indica*) exhibits potent antibacterial and antifungal activities through azadirachtin and nimbidin compounds, making it effective against acne-causing bacteria and dermatophytes [13]. Turmeric (*Curcuma longa*) curcumin displays antimicrobial properties while simultaneously reducing inflammatory responses in infected skin.

Anti-aging and Skin Regeneration Effects

Botanical ingredients counteract intrinsic and extrinsic aging through multiple mechanisms including collagen stimulation, elastin preservation, and cellular renewal. Ginseng (*Panax ginseng*) ginsenosides promote fibroblast proliferation, increase hyaluronic acid synthesis, and reduce wrinkle formation [14]. Retinol-like compounds from rosehip (*Rosa canina*) stimulate collagen production and accelerate cell turnover, improving skin texture and reducing fine lines [15]. Centella asiatica triterpenes (asiaticoside, madecassoside) enhance collagen synthesis and improve skin firmness through SMAD signalling pathway activation.

Cosmetic Applications of Medicinal Plants and Herbs

Skin Whitening and Brightening Agents

Plant-based tyrosinase inhibitors offer safe alternatives to synthetic depigmenting agents. Licorice (*Glycyrrhiza glabra*) glabridin inhibits tyrosinase activity and reduces melanin production without cytotoxicity [16]. Mulberry (*Morus alba*) extract contains oxyresveratrol, effectively suppressing melanogenesis and treating hyperpigmentation [17]. Kojic acid, derived from *Aspergillus oryzae*, chelates copper ions in tyrosinase active sites, preventing melanin formation.

Hair Care: Strengthening, Growth Promotion, and Scalp Health

Botanical extracts address various hair concerns through nutritive and stimulatory properties. Saw palmetto (*Serenoa repens*) inhibits 5 α -reductase, reducing DHT-mediated hair loss in androgenetic alopecia [18]. Rosemary (*Rosmarinus officinalis*) improves scalp circulation and demonstrates efficacy comparable to minoxidil in promoting hair growth. Hibiscus (*Hibiscus rosa-sinensis*) amino acids and vitamins strengthen hair shafts, prevent premature graying, and condition scalp.

Natural Moisturizers and Emollients

Plant oils and extracts provide superior hydration through occlusive, humectant, and emollient properties. Hyaluronic acid from botanical fermentation attracts moisture, maintaining skin hydration and plumpness [19]. Shea butter (*Vitellaria paradoxa*) triterpenes and fatty acids restore skin barrier function and provide long-lasting moisturization [20]. Avocado (*Persea americana*) oil penetrates deep into skin layers, delivering essential fatty acids and vitamins that enhance barrier integrity [21].

Treatment of Acne and Other Skin Disorders

Medicinal plants offer comprehensive acne management through antibacterial, anti-inflammatory, and sebum-regulating effects. Witch hazel (*Hamamelis virginiana*) tannins provide astringent properties, reducing sebum production and pore size [22]. Willow bark (*Salix alba*) salicin acts as a natural beta-hydroxy acid, exfoliating pores and preventing comedone formation [23]. Tea tree oil demonstrates efficacy comparable to benzoyl peroxide in reducing acne lesions with fewer side effects.

Key Bioactive Compounds Contributing to Cosmeceutical Effects

Polyphenols and Flavonoids

Polyphenols represent the most extensively studied phytochemicals in cosmeceutical applications, exhibiting potent antioxidant, anti-inflammatory, and photoprotective properties. Catechins from green tea, particularly EGCG, neutralize free radicals and inhibit UV-induced matrix metalloproteinase expression, preventing collagen degradation [24]. Resveratrol, a stilbenoid polyphenol from grapes and berries, activates sirtuins and enhances cellular longevity while protecting against oxidative stress [25]. Flavonoids such as quercetin and kaempferol demonstrate anti-aging effects through modulation of NF- κ B signaling and reduction of inflammatory mediators.

Essential Oils and Terpenoids

Essential oils contain volatile terpenoids that penetrate skin barriers effectively, delivering therapeutic benefits. Limonene exhibits anti-inflammatory and wound-healing properties while enhancing transdermal absorption of other compounds [26]. Linalool and geraniol demonstrate antimicrobial activity against skin pathogens and provide soothing aromatherapeutic effects [27]. Triterpenes such as ursolic acid and oleanolic acid stimulate collagen synthesis, inhibit elastase activity, and promote skin regeneration.

Alkaloids and Glycosides

Plant alkaloids offer diverse cosmeceutical benefits through receptor-mediated mechanisms. Caffeine constricts blood vessels, reduces puffiness, and demonstrates lipolytic activity in cellulite treatment [28]. Berberine exhibits antimicrobial and anti-inflammatory properties beneficial for acne management [29]. Glycosides such as arbutin inhibit tyrosinase competitively, reducing melanin synthesis and treating hyperpigmentation [30]. Saponins from ginseng enhance skin permeability and exhibit immunomodulatory effects [31].

Vitamins and Minerals

Plant-derived vitamins function as essential cofactors in skin metabolism and protection. Vitamin C (ascorbic acid) serves as a cofactor for prolyl hydroxylase in collagen synthesis while neutralizing ROS [32]. Vitamin E (tocopherols) protects cell membranes from lipid peroxidation and works synergistically with vitamin C [32]. Phyto mineral complexes provide trace elements like zinc and selenium necessary for antioxidant enzyme function and wound healing [33].

Safety, Efficacy, and Regulatory Aspects

Toxicity and Allergenicity Considerations

Despite natural origins, botanical cosmeceuticals pose potential safety concerns including contact dermatitis, phototoxicity, and systemic toxicity. Essential oils, particularly tea tree and lavender, may cause allergic reactions in sensitized individuals [34]. Furanocoumarins in citrus oils induce phototoxic responses upon UV exposure [35]. Comprehensive safety assessments including patch testing, repeated insult patch tests, and photo safety evaluations are essential before commercialization [36].

Clinical Evidence and Validation of Medicinal Plant Cosmeceuticals

Rigorous clinical validation remains crucial for substantiating cosmeceutical claims. Randomized controlled trials demonstrate green tea polyphenols significantly reduce photoaging signs including wrinkles and hyperpigmentation [37]. Double-blind studies confirm niacinamide effectively reduces melanin transfer and improves skin barrier function [38]. However, many botanical ingredients lack sufficient clinical evidence, necessitating standardized methodologies for efficacy assessment [39].

Regulatory Framework and Quality Control

Cosmeceutical regulation varies globally, creating market complexities. The FDA does not recognize cosmeceuticals as a distinct category, classifying products as either cosmetics or drugs based on intended use [40]. European Union regulations require safety assessments and good manufacturing practices for cosmetic ingredients [41].

Quality control measures including phytochemical standardization, contaminant testing, and stability studies ensure product consistency and safety [42].

Challenges and Future Perspectives

Standardization and Formulation Challenges

Botanical extract standardization presents significant challenges due to phytochemical variability influenced by cultivation conditions, harvesting time, and extraction methods (Heinrich et al., 2004). Establishing marker compounds and implementing fingerprinting techniques ensure batch-to-batch consistency [43]. Formulation challenges include low water solubility, chemical instability, and poor skin penetration of many phytoactives [44].

Advances in Phytochemical Extraction and Delivery Systems

Novel extraction technologies including supercritical fluid extraction, ultrasound-assisted extraction, and enzyme-assisted extraction enhance bioactive yield while preserving compound integrity [45]. Advanced delivery systems such as niosomes, phytosomes, and solid lipid nanoparticles improve phytochemical stability, penetration, and sustained release [46]. Microencapsulation protects sensitive compounds from degradation and enables controlled release [47].

Potential for Novel Medicinal Plants in Cosmeceutical Development

Bioprospecting unexplored botanical resources offers tremendous potential for discovering novel cosmeceutical ingredients. Marine botanicals like seaweeds contain unique polysaccharides and phlorotannins with superior antioxidant properties [48]. Adaptogenic herbs including rhodiola and ashwagandha demonstrate stress-protective effects on skin (Wiegant et al., 2009). Integration of systems biology, metabolomics, and artificial intelligence accelerates identification and optimization of botanical cosmeceuticals [49].

Conclusion

Plant-based cosmeceuticals offer scientifically validated therapeutic benefits through diverse bioactive compounds. Continued research in standardization, delivery systems, and clinical validation will advance their integration into evidence-based dermatological practice, establishing botanical ingredients as cornerstone elements in modern skincare formulations.

References

- Shah, P., Banerjee, S., Singh, A., Kulhari, H., & Anand Saharan, V. (2025). From traditional to modern medicine: the role of herbs and phytoconstituents in pharmaceuticals, nutraceuticals, and cosmetics. *Formulating Pharma-, Nutra-, and Cosmeceutical Products from Herbal Substances: Dosage Forms and Delivery Systems*, 3-73.
- Ahmed, I. A., Dia, K., Alhazmi, A. L. A., & Saeid, M. J. E. (2025). Medicinal Plants in Cosmeceuticals: Bridging Ethnopharmacology and Modern Dermatology. *International Journal of Biotechnology and Biomedicine (IJBB)*, 2(3), 45-58.
- Nadeeshani Dilhara Gamage, D. G., Dharmadasa, R. M., Chandana Abeysinghe, D., Saman Wijesekara, R. G., Prathapasinghe, G. A., & Someya, T. (2022). Global perspective of plant-based cosmetic industry and possible contribution of Sri Lanka to the development of herbal cosmetics. *Evidence-Based Complementary and Alternative Medicine*, 2022(1), 9940548.

4. Akihisa, T., Kojima, N., Katoh, N., Ichimura, Y., Suzuki, H., Fukatsu, M., ... & Yasukawa, K. (2010). Triterpene alcohol and sterol ferulates from shea nut and their anti-inflammatory and antitumor-promoting effects. *Journal of Natural Products*, 73(11), 1812-1817.
5. Alzohairy, M. A. (2016). Therapeutics role of *Azadirachta indica* (Neem) and their active constituents in diseases prevention and treatment. *Evidence-Based Complementary and Alternative Medicine*, 2016, 7382506.
6. Lall, N., Mahomoodally, M. F., Esposito, D., Steenkamp, V., Zengin, G., Steyn, A., & Oosthuizen, C. B. (2020). Cosmeceuticals from medicinal plants. *Frontiers in Pharmacology*, 11, 1149.
7. Ola, M., Bhaskar, R., Patil, P. S., Badgular, S. V., & Haswani, N. G. (2025). Formulation and Delivery Issues for Active Ingredients of Herbal Medicines, Nutraceuticals, and Cosmetics: Comparisons to Small-Molecule Drugs. *Formulating Pharma-, Nutra-, and Cosmeceutical Products from Herbal Substances: Dosage Forms and Delivery Systems*, 163-191.
8. Basketter, D. A., Clewell, H., Kimber, I., Rossi, A., Blauboer, B., Burrier, R., ... & Hasiwa, M. (2008). A roadmap for the development of alternative (non-animal) methods for systemic toxicity testing. *ALTEX*, 29(1), 3-91.
9. Baxter, R. A. (2008). Anti-aging properties of resveratrol: Review and report of a potent new antioxidant skin care formulation. *Journal of Cosmetic Dermatology*, 7(1), 2-7.
10. Bissett, D. L., Oblong, J. E., & Berge, C. A. (2004). Niacinamide: A B vitamin that improves aging facial skin appearance. *Dermatologic Surgery*, 31(s1), 860-866.
11. Khan, S., Ghawte, S. A., Ghawte, S., Mulla, G., & Shaikh, N. (2025). Herbal Plants and Their Applications in Cosmeceuticals: A Comprehensive Review. *Plant Science Review* DOI: <https://doi.org/10.51470/PSR.2025.06.02.44>
12. Bonifácio, B. V., Silva, P. B., Ramos, M. A., Negri, K. M., Bauab, T. M., & Chorilli, M. (2014). Nanotechnology-based drug delivery systems and herbal medicines. *International Journal of Nanomedicine*, 9, 1-15.
13. Vitalone, A., D'Andrea, L., Di Sotto, A., Caruso, A., & Parente, R. (2025). Official plants as new frontiers of cosmetic ingredients. *Cosmetics*, 12(4), 140.
14. Carson, C. F., & Riley, T. V. (1995). Antimicrobial activity of the major components of the essential oil of *Melaleuca alternifolia*. *Journal of Applied Bacteriology*, 78(3), 264-269.
15. Carson, C. F., Hammer, K. A., & Riley, T. V. (2006). *Melaleuca alternifolia* (tea tree) oil: A review of antimicrobial and other medicinal properties. *Clinical Microbiology Reviews*, 19(1), 50-62.
16. Chanchal, D., & Swarnlata, S. (2008). Novel approaches in herbal cosmetics. *Journal of Cosmetic Dermatology*, 7(2), 89-95.
17. Vartak, A., Sonawane, S., Alim, H., Patel, N., Hamrouni, L., Khan, J., & Ali, A. (2022). Medicinal and aromatic plants in the cosmetics industry. In *Medicinal and Aromatic Plants of India Vol. 1* (pp. 341-364). Cham: Springer International Publishing.
18. Choi, K. T. (2018). Botanical characteristics, pharmacological effects and medicinal components of Korean *Panax ginseng* C A Meyer. *Acta Pharmacologica Sinica*, 29(9), 1109-1118.
19. Comunian, T. A., & Favaro-Trindade, C. S. (2016). Microencapsulation using biopolymers as an alternative to produce food enhanced with phytosterols and omega-3 fatty acids. *Food Hydrocolloids*, 61, 442-457.
20. Gamage, D. G. N. D., Dharmadasa, R. M., Abeysinghe, D. C., Wijesekara, R. G. S., Prathapasinghe, G. A., & Someya, T. (2021). Ethnopharmacological survey on medicinal plants used for cosmetic treatments in traditional and ayurveda systems of medicine in Sri Lanka. *Evidence-Based Complementary and Alternative Medicine*, 2021(1), 5599654.
21. Draelos, Z. D. (2009). Cosmeceuticals: Undefined, unclassified, and unregulated. *Clinics in Dermatology*, 27(5), 431-434.
22. Dugrand-Judek, A., Olry, A., Hehn, A., Costantino, G., Ollitrault, P., Froelicher, Y., & Bourgaud, F. (2015). The distribution of coumarins and furanocoumarins in citrus species closely matches citrus phylogeny. *Phytochemistry*, 110, 65-74.
23. Hammer, K. A., Carson, C. F., Riley, T. V., & Nielsen, J. B. (2012). A review of the toxicity of *Melaleuca alternifolia* (tea tree) oil. *Food and Chemical Toxicology*, 44(5), 616-625.
24. Heinrich, M., Barnes, J., Gibbons, S., & Williamson, E. M. (2004). *Fundamentals of pharmacognosy and phytotherapy*. Churchill Livingstone.
25. Heinrich, U., Moore, C. E., De Spirt, S., Tronnier, H., & Stahl, W. (2011). Green tea polyphenols provide photoprotection, increase microcirculation, and modulate skin properties of women. *Journal of Nutrition*, 141(6), 1202-1208.
26. Herman, A., & Herman, A. P. (2013). Caffeine's mechanisms of action and its cosmetic use. *Skin Pharmacology and Physiology*, 26(1), 8-14.
27. Ilyasoglu, H., & Gultekin, M. S. (2013). A mini review on rosehip fruit: A detailed scenario on processing and utilization of rosehip. *African Journal of Traditional, Complementary and Alternative Medicines*, 10(6), 549-553.
28. Jäger, S., Trojan, H., Kopp, T., Laszczyk, M. N., & Scheffler, A. (2009). Pentacyclic triterpene distribution in various plants-rich sources for a new group of multi-potent plant extracts. *Molecules*, 14(6), 2016-2031.
29. Katiyar, S. K., Ahmad, N., & Mukhtar, H. (2011). Green tea and skin. *Archives of Dermatology*, 136(8), 989-994.
30. Kerschner, M., Buntrock, H., Bertram, A., Küster, W., & Meurer, M. (2011). Stability testing of cosmetic emulsions. *Skin Pharmacology and Physiology*, 4(1), 77-81.
31. Kligman, D. E. (2000). Cosmeceuticals. *Dermatologic Clinics*, 18(4), 609-615.
32. Lee, K. T., Lee, K. S., Jeong, J. H., Jo, B. K., Heo, M. Y., & Kim, H. P. (2002). Inhibitory effects of *Ramulus mori* extracts on melanogenesis. *Journal of Cosmetic Science*, 53(6), 343-348.
33. Leyden, J. J., Shergill, B., Micali, G., Downie, J., & Wallo, W. (2011). Natural options for the management of hyperpigmentation. *Journal of the European Academy of Dermatology and Venereology*, 25(10), 1140-1145.
34. Leyva-López, N., Gutierrez-Grijalva, E. P., Ambriz-Perez, D. L., & Heredia, J. B. (2016). Flavonoids as cytokine modulators: A possible therapy for inflammation-related diseases. *International Journal of Molecular Sciences*, 17(6), 921.
35. Man, M. Q., Yang, B., & Elias, P. M. (2008). Benefits of hesperidin for cutaneous functions. *Evidence-Based Complementary and Alternative Medicine*, 6(S1), 23-28.
36. Masum, M. N., Yamauchi, K., & Mitsunaga, T. (2019). Tyrosinase inhibitors from natural and synthetic sources as skin-lightening agents. *Reviews in Agricultural Science*, 7, 41-58.
37. McKay, D. L., & Blumberg, J. B. (2006). A review of the bioactivity and potential health benefits of chamomile tea. *Phytotherapy Research*, 20(7), 519-530.
38. Mukherjee, P. K., & Houghton, P. J. (2009). *Evaluation of herbal medicinal products*. Pharmaceutical Press.
39. Mukherjee, P. K., Maity, N., Nema, N. K., & Sarkar, B. K. (2011). Bioactive compounds from natural resources against skin aging. *Phytomedicine*, 19(1), 64-73.
40. Mukherjee, P. K., Biswas, R., Sharma, A., Banerjee, S., Biswas, S., & Katiyar, C. K. (2014). Validation of medicinal herbs for anti-tyrosinase potential. *Journal of Herbal Medicine*, 4(4), 171-193.
41. Murugusundram, S. (2009). *Serenoa repens*: Does it have any role in the management of androgenetic alopecia? *Journal of Cutaneous and Aesthetic Surgery*, 2(1), 31-32.

42. Nichols, J. A., & Katiyar, S. K. (2010). Skin photoprotection by natural polyphenols: Anti-inflammatory, antioxidant and DNA repair mechanisms. *Archives of Dermatological Research*, 302(2), 71-83.
43. Panahi, Y., Taghizadeh, M., Marzony, E. T., & Sahebkar, A. (2015). Rosemary oil vs minoxidil 2% for the treatment of androgenetic alopecia: A randomized comparative trial. *Skinmed*, 13(1), 15-21.
44. Papakonstantinou, E., Roth, M., & Karakiulakis, G. (2012). Hyaluronic acid: A key molecule in skin aging. *Dermato-Endocrinology*, 4(3), 253-258.
45. Pillaiyar, T., Manickam, M., & Namasivayam, V. (2017). Skin whitening agents: Medicinal chemistry perspective of tyrosinase inhibitors. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 32(1), 403-425.
46. Poljšak, B., & Dahmane, R. (2012). Free radicals and extrinsic skin aging. *Dermatology Research and Practice*, 2012, 135206.
47. Pullar, J. M., Carr, A. C., & Vissers, M. C. (2017). The roles of vitamin C in skin health. *Nutrients*, 9(8), 866.
48. Rinnerthaler, M., Bischof, J., Streubel, M. K., Trost, A., & Richter, K. (2015). Oxidative stress in aging human skin. *Biomolecules*, 5(2), 545-589.
49. Rostan, E. F., DeBuys, H. V., Madey, D. L., & Pinnell, S. R. (2002). Evidence supporting zinc as an important antioxidant for skin. *International Journal of Dermatology*, 41(9), 606-611.